

Berichte des  
Bundesamtes für Seeschiffahrt und Hydrographie  
Nr. 23

**The Subsurface Eastern Boundary Current of  
the North Atlantic between 32°N and 53°N**

---

**Data Report 1983–1995**

**Author: Holger Klein**




Hamburg und Rostock 2000



## **Contents**

Introduction	1
Data processing	4
Moorings F1, F2, F4	7
Moorings G1 –G5	41
Mooring L01	97
Mooring AvH	105
Mooring K1	127
Mooring WH	139
Mooring P3	147
Mooring CS	161
Moorings AL1, AL2	179
Mooring R	211
Mooring M3	227



## I. Introduction

This data report presents current data collected by the Bundesamt für Seeschiffahrt und Hydrographie (BSH, Federal Maritime and Hydrographic Agency) between 1983 and 1995 along the European and Northwest African continental slope between 32°N and 53°N.

The mooring positions are located along the coast of the Iberian Peninsula, the northern part of the Moroccan coast (M3, R), in the Great and Little Sole Bank area (G-moorings), and west of the Irish Coast close to the Porcupine Bank (F-moorings).

The deployment periods vary between 15 and 414 days. All current metres were Aanderaa RCM 4, 5, 7, or 8. The sampling interval for the long-term moorings was 60 minutes, for the shorter deployment periods (< 6 months) the sampling interval varies between 5 and 30 minutes.

Some of the data have been discussed in previous reports<sup>1,2</sup>, but this report includes all BSH current data relevant to the subsurface Eastern Boundary Current north of 32°N. Table 1 and the map in Figure 1 give the mooring positions together with deployment time, water depth, and the number of current meter records in the respective mooring.

For each mooring there is a table with mooring data followed by a table with low-passed current statistics and a progressive vector diagram (hereafter PVD) of all current meters in the mooring. For each record there is a time series plot (low-passed), an energy density spectra for the zonal and meridional current component, and an individual PVD. For the current statistics and the time series a Gaussian low-pass over 48 hours was applied to remove short-term fluctuations such as tidal and inertial movements.

The PVDs were processed from unfiltered data. The start position of the PVDs is marked by a big cross (+). The spacing of the time marks (x) depends upon the record length and is given in the text of the diagram.

The spectra are Blackman-Tukey power spectra calculated with the Singular-Spectrum-Analysis Toolkit<sup>3</sup>.

<sup>1</sup> Mittelstaedt, E., J. Meincke and Holger Klein, 1996: WOCE-Current measurements: The ACM8 array. Data Report. Berichte des Bundesamtes für Seeschiffahrt und Hydrographie Nr.8, 150pp.

<sup>2</sup> Hagen, E. et al., 1994: Hydrographische Untersuchungen im Ostrandstromsystem vor Portugal und Marokko 1991 - 1992. Berichte des Bundesamtes für Seeschiffahrt und Hydrographie Nr.2, 50pp.

<sup>3</sup> The Singular-Spectrum\_analysis Toolkit User's Guide. Department of Atmospheric Sciences, University of California, Los Angeles, 1995, 50pp.




Figure 1: Mooring positions. Isobaths are given for 200 m and 2000 m depth.

**Table1: Mooring positions**

mooring idendifier	position	deployment time	water depth	nc
-	-	-	m	-
F1	52°20.4 'N	16°20.8 'W	19.09.91 - 30.08.92	3510 3
F2	52°22.7 'N	16°21.7 'W	30.08.92 - 11.09.93	3481 5
F4	52°23.2 'N	15°28.5 'W	23.05.94 - 31.05.95	2554 4
G1	48°55.8 'N	12°55.8 'W	16.06.93 - 5.08.94	2540 3
G2	48°50.8 'N	13°59.5 'W	17.06.93 - 15.02.94	4530 2
G3	50°17.4 'N	11°53.7 'W	13.06.94 - 3.07.94	2100 5
G4	47°59.3 'N	9° 2.2 'W	14.06.94 - 8.07.94	2500 6
G5	48°43.9 'N	13° 8.0 'W	6.08.95 - 21.08.95	3900 5
L01	42°55.54'N	9°50.88'W	14.03.91 - 19.01.92	1945 1
AvH	42°32.80'N	14°28.60'W	1.10.89 - 9.08.90	5165 8
K1	40°47.4 'N	9°06.2 'W	19.03.91 - 13.04.91	105 3
WH	40°13.3 'N	10°15.7 'W	15.08.90 - 27.08.91	3900 2
P3	40° 1.8 'N	9°45.2 'W	30.07.83 - 14.01.84	910 4
CS	36°37.1 'N	9°22.0 'W	28.03.92 - 19.05.92	1960 6
AL1	36°14.16'N	8°36.12'W	20.10.91 - 14.01.92	2036 6
AL2	36°14.20'N	8°35.10'W	14.01.92 - 19.05.92	1998 6
R	34°29.10'N	7°50.8 'W	20.10.91 - 18.05.92	1995 5
M3	32°35.72'N	10°49.18'W	18.01.92 - 25.05.92	3979 1

nc = number of current meter records per deployment

## II. Data processing

The Aanderaa RCM raw data are converted to physical data using individual calibration data for each current meter. The BSH calibrates its current meters regularly performing tests for rotor speed and compass deviation.

Current statistics are calculated from data smoothed over 48 hours with a Gaussian low-pass filter to remove tidal currents. The mean values of the zonal ( $u$ ) and meridional ( $v$ ) components of current vector  $V$  are given together with the 95% confidence interval assuming a Gaussian normal distribution:

$$\text{mean value} \pm 1.96 \frac{\sqrt{u_i' u_i'}}{\sqrt{n-1}}, n = \text{number of cycles}$$

The Lagrangian statistics were calculated according to Taylor (1921)<sup>4</sup>.

### Symbols and abbreviations used in the tables:

dir mean current direction

$$dir(\bar{u}, \bar{v}) = atan2(\bar{u}, \bar{v}) \times \frac{180}{\pi}$$

$k_M$  mean kinetic energy

$$k_M = \frac{1}{2}(\bar{u}^2 + \bar{v}^2)$$

$k_E$  eddy kinetic energy

$$k_E = \frac{1}{2}(\bar{u}'\bar{u}' + \bar{v}'\bar{v}')$$

$K_{x,y}$  eddy diffusivity

$$K_i = \bar{u}_i' \bar{u}_i' T_i$$

$L_{x,y}$  Lagrangian length scale

$$L_i = \sqrt{\bar{u}_i' \bar{u}_i'} T_i$$

$mag$  magnitude of current vector  $V$

RCM recording current meter

<sup>4</sup> Taylor, G.I., 1921: Diffusion by continuous movements.  
Proc. London Math. Soc., A20, 196-211.

si	sampling interval
sd	sampling depth
$T_{x,y}$	Lagrangian integral time scale
	$T_i = \int_0^{\infty} R_i(\tau) d\tau \quad , \text{with} \quad R_i = \frac{1}{u_i' u_i T_{\max}} \int_0^{T_{\max}} u_i'(t) u_i'(t + \tau) dt$
$\bar{u}, \bar{v}$	zonal and meridional component of $V$
$\overline{u'u'}$	zonal variance
$\overline{u'v'}$	covariance
$\overline{v'v'}$	meridional variance
$V$	mean current vector



F1, F2, F4



### Mooring Data F1, F2, and F4


---

current meter	sampl. depth	mab	record no.	sampling time		
-	m	m	-	UTC		
<b>F1: 19.9.91 - 30.8.92, 52° 20.4'N, 16° 20.8'W, si = 60'</b>						
RCM5	6784	210	3300	92067	19.9.91 14:00	- 30.8.92 6:00
RCM5	6789	3010	500	92063	19.9.91 14:00	- 30.8.92 6:00
RCM5	6822	3460	50	92062	19.9.91 14:00	- 19.2.92 12:00
<b>F2: 30.8.92 - 11.9.93, 52° 22.7'N, 16° 21.7'W, si = 60'</b>						
RCM4	996	181	3300	93049	30.8.92 18:00	- 5.8.93 8:00
RCM5	6890	781	2700	93047	30.8.92 18:00	- 5.9.93 7:00
RCM5	6969	1981	1500	93046	30.8.92 18:00	- 11.9.93 17:00
RCM8	10082	2981	200	93045	30.8.92 18:00	- 4.9.93 1:00
RCM8	9358	3431	50	93044	30.8.92 18:00	- 5.8.93 23:00
<b>F4: 23.5.94 - 31.5.95, 52° 23.2'N, 15° 28.5'W, si = 60'</b>						
RCM5	6968	853	1701	95023	23.5.94 10:00	- 31.5.95 7:00
RCM5	6835	1255	1299	95024	23.5.94 10:00	- 31.5.95 7:00
RCM5	6969	2057	497	95025	23.5.94 10:00	- 31.5.95 7:00
RCM5	6890	2504	50	95026	23.5.94 10:00	- 19.5.95 16:00

mab = metres above bottom, si = sampling interval

Current statistics (low-passed data)


Id	sampl. depth	rec. length	$\bar{U}$	$\bar{V}$	mag $\mathbf{v}$	$\frac{\bar{u}'\bar{u}'}{\bar{v}'\bar{v}'}$	$\frac{\bar{u}'\bar{v}'}{\bar{v}'\bar{v}'}$	KE			Tx			Lx			Kx			dir		
								m	days	cm/s	cm/s	cm <sup>2</sup> /s <sup>2</sup>	cm <sup>2</sup> /s <sup>2</sup>	cm <sup>2</sup> /s <sup>2</sup>	days	km	cm <sup>2</sup> /s <sup>2</sup>	km	km	km	km	km
<b>F1:</b>	210	345	-2.8±0.26	20.2	286.8	-64.0	214.2	21.4				313.6	531.2				246					
			-1.3±0.36	3.1	141.6		5.0	9.1				93.4	111.1									
	3010	345	-0.2±0.06	4.9	8.7	-0.5	9.0	1.9				4.9	1.4				249					
			-0.1±0.07	0.3	9.3		<0.1	1.8				4.7	1.4									
	3460	152	-0.4±0.18	6.3	30.1	-0.4	18.1	7.9				37.6	20.6				214					
			-0.5±0.08	0.7	6.1		0.2	1.8				3.8	0.9									
<b>F2:</b>	181	340	-2.5±0.19	13.9	80.6	13.7	76.8	7.5				58.5	52.5				240					
			-1.4±0.19	2.8	73.0		4.1	21.8				161.3	137.9									
	781	370	-1.2±0.15	10.5	50.5	12.8	49.1	10.0				61.3	43.5				228					
			-1.1±0.14	1.6	47.7		1.3	31.9				190.6	131.6									
	1981	377	-2.2±0.14	6.8	24.5	-0.2	21.2	8.6				36.7	18.2				264					
			-0.2±0.05	2.2	18.1		2.5	25.5				93.8	39.9									
<b>F3:</b>	2981	369	-0.2±0.08	5.0	7.4	1.6	6.4	2.1				4.9	1.3				195					
			-0.8±0.06	0.9	5.5		0.4	3.3				6.8	1.6									
	3431	340	-0.7±0.10	6.9	20.0	2.5	13.5	1.8				6.8	3.0				233					
			-0.5±0.08	0.9	13.5		0.4	2.6				8.3	3.0									
<b>F4:</b>	853	371	0.3±0.08	7.1	16.2	8.4	22.3	26.6				92.5	37.2				174					
			-2.4±0.08	2.5	28.5		3.0	20.0				92.4	49.3									
	1255	371	-0.8±0.09	8.3	19.8	12.0	32.1	24.1				92.5	41.1				189					
			-4.8±0.14	4.8	44.3		11.6	15.5				89.1	59.3									
	2057	371	-0.6±0.06	7.4	7.2	-0.7	20.6	7.7				17.8	4.8				199					
			-1.8±0.12	1.9	34.1		1.7	11.8				59.5	34.7									
<b>F5:</b>	2504	359	-0.8±0.02	8.1	1.1	1.0	5.0	1.9				4.7	0.2				345					
			2.8±0.06	2.9	8.9		4.3	1.8				4.7	1.4									



Time marks (x) are given every 720 hours

- A** o92067; sd= 210 m; 19. 9.1991 14:00 – 30. 8.1992 6:00 (UTC)
- B** o92063; sd= 3010 m; 19. 9.1991 14:00 – 30. 8.1992 6:00 (UTC)
- C** o92062; sd= 3460 m; 19. 9.1991 14:00 – 19. 2.1992 12:00 (UTC)


F1, record no. = 092067, sampling depth = 210 m





F1

record no: a92067, sampling interval = 60.0 minutes

u-component (zonal)




v-component (meridional)



19. 9.1991 14:00 – 30. 8.1992 6:00 (UTC)

Time marks (x) are given every 720 hours


F1, record no. = a92063, sampling depth = 3010 m





F1

record no: a92063, sampling interval = 60.0 minutes

u-component (zonal)




v-component (meridional)



19. 9.1991 14:00 – 30. 8.1992 6:00 (UTC)

Time marks (x) are given every 720 hours


F1, record no. = a92062, sampling depth = 3460 m





F1

record no: a92062, sampling interval = 60.0 minutes

u-component (zonal)




v-component (meridional)



19. 9.1991 14:00 – 19. 2.1992 12:00 (UTC)


Time marks (x) are given every 720 hours





- A** o93049; sd= 181 m; 30. 8.1992 18:00 – 5. 8.1993 8:00 (UTC)
- B** o93047; sd= 781 m; 30. 8.1992 18:00 – 5. 9.1993 7:00 (UTC)
- C** o93046; sd= 1981 m; 30. 8.1992 18:00 – 11. 9.1993 17:00 (UTC)
- D** o93045; sd= 2981 m; 30. 8.1992 18:00 – 4. 9.1993 1:00 (UTC)
- E** o93044; sd= 3431 m; 30. 8.1992 18:00 – 5. 8.1993 23:00 (UTC)


F2, record no. = 093049, sampling depth = 181 m





F2

record no: a93049, sampling interval = 60.0 minutes

u-component (zonal)




v-component (meridional)



30. 8.1992 18:00 – 5. 8.1993 8:00 (UTC)

Time marks (x) are given every 720 hours


F2, record no. = a93047, sampling depth = 781 m





F2

record no: a93047, sampling interval = 60.0 minutes

u-component (zonal)




v-component (meridional)



30. 8.1992 18:00 – 5. 9.1993 7:00 (UTC)



Time marks (x) are given every 720 hours

F2, record no. = a93046, sampling depth = 1981 m



F2


record no: a93046, sampling interval = 60.0 minutes



30. 8.1992 18:00 – 11. 9.1993 17:00 (UTC)

Time marks (x) are given every 720 hours


F2, record no. = 093045, sampling depth = 2981 m





F2

record no: a93045, sampling interval = 60.0 minutes

u-component (zonal)




v-component (meridional)



30. 8.1992 18:00 – 4. 9.1993 1:00 (UTC)

Time marks (x) are given every 720 hours


F2, record no. = a93044, sampling depth = 3431 m





F2

record no: a93044, sampling interval = 60.0 minutes

u-component (zonal)




v-component (meridional)



30. 8.1992 18:00 – 5. 8.1993 23:00 (UTC)

Time marks (x) are given every 720 hours






**A** o95023; sd= 853 m; 23. 5.1994 10:00 – 31. 5.1995 7:00 (UTC)

**B** o95024; sd= 1255 m; 23. 5.1994 10:00 – 31. 5.1995 7:00 (UTC)

**C** o95025; sd= 2057 m; 23. 5.1994 10:00 – 31. 5.1995 7:00 (UTC)

**D** o95026; sd= 2504 m; 23. 5.1994 10:00 – 19. 5.1995 16:00 (UTC)


F4, record no. = 095023, sampling depth = 853 m





F4

record no: a95023, sampling interval = 60.0 minutes

u-component (zonal)




v-component (meridional)



23. 5.1994 10:00 – 31. 5.1995 7:00 (UTC)

Time marks (x) are given every 720 hours


F4, record no. = a95024, sampling depth = 1255 m





F4

record no: a95024, sampling interval = 60.0 minutes

u-component (zonal)




v-component (meridional)



23. 5.1994 10:00 – 31. 5.1995 7:00 (UTC)

Time marks (x) are given every 720 hours


F4, record no. = a95025, sampling depth = 2057 m





F4

record no: a95025, sampling interval = 60.0 minutes

u-component (zonal)




v-component (meridional)



23. 5.1994 10:00 – 31. 5.1995 7:00 (UTC)

Time marks (x) are given every 720 hours


F4, record no. = a95026, sampling depth = 2504 m





F4

record no: a95026, sampling interval = 60.0 minutes

u-component (zonal)



v-component (meridional)



23. 5.1994 10:00 – 19. 5.1995 16:00 (UTC)

Time marks (x) are given every 720 hours



G1 - G5



### Mooring Data G1, G2, G3, G4, and G5

current meter	sampl. depth	mab	record no.		sampling time					
-	m	m	-		UTC					
<b>G1: 16.6.93 - 5.8.94, 48° 55.8'N, 12° 55.8'W, si = 60'</b>										
RCM5 6786 632 1908 95047 16.6.93 21:00 - 12.7.94 23:00										
RCM5	6787	1035	1505	95048	16.6.93	22:00	- 5.8.94 14:00			
RCM5	6976	1838	702	95049	16.6.93	22:00	- 11.7.94 07:00			
<b>G2: 17.6.93 - 15.2.94, 48° 50.8'N, 13° 59.5'W, si = 60'</b>										
RCM8	10588	370	4160	95051	17.6.93	12:00	- 4.12.93 04:00			
RCM8	10586	977	3553	95053	17.6.93	13:00	- 15.2.94 02:00			
<b>G3: 13.6.94 - 3.7.94, 50° 17.4'N, 11° 53.7'W, si = 10'</b>										
RCM4	4731	367	1733	94037	13.6.94	09:00	- 3.7.94 05:50			
RCM4	4729	579	1521	94038	13.6.94	09:00	- 3.7.94 05:50			
RCM4	4364	881	1219	94039	13.6.94	09:00	- 3.7.94 05:50			
RCM4	6107	1383	717	94040	13.6.94	09:00	- 3.7.94 05:50			
RCM8	10649	2085	15	94041	13.6.94	09:00	- 3.7.94 05:50			
<b>G4: 14.6.94 - 8.7.94, 47° 59.3'N, 9° 2.2'W, si = 10'</b>										
RCM4	7752	435	2065	94036	14.6.94	12:20	- 8.7.94 07:50			
RCM4	4742	587	1913	94035	14.6.94	12:20	- 8.7.94 07:50			
RCM4	7750	739	1761	94034	14.6.94	12:20	- 8.7.94 07:50			
RCM4	7751	941	1559	94033	14.6.94	12:20	- 8.7.94 07:50			
RCM8	10650	2485	15	94031	14.6.94	12:20	- 8.7.94 07:50			
<b>G5: 6.8.95 - 21.8.95, 48° 43.9'N, 13° 8.0'W, si = 5'</b>										
RCM7	11556	140	3760	95036	6.8.95	15:15	- 21.8.95 14:10			
RCM8	10684	340	3560	95037	6.8.95	15:15	- 21.8.95 14:10			
RCM8	9385	540	3360	95038	6.8.95	15:15	- 21.8.95 14:15			
RCM8	10649	1140	2760	95039	6.8.95	15:15	- 21.8.95 14:10			
RCM8	9384	2140	1760	95040	6.8.95	15:15	- 21.8.95 14:10			

mab = metres above bottom, si = sampling interval

Current statistics (low-passed data)


Id	sampl. depth	rec. length	$\bar{U}$ $\bar{V}$	mag $\mathbf{v}$	$\frac{\bar{U}'\bar{U}'}{\bar{V}'\bar{V}'}$	$\bar{U}'\bar{V}'$	$k_E$ $k_N$	$T_x$ $T_y$	$L_x$ $L_y$	$K_x$ $K_y$	dir			
	m	days	cm/s	cm/s	$\text{cm}^2/\text{s}^2$	$\text{cm}^2/\text{s}^2$	$\text{cm}^2/\text{s}^2$	days	km	$\text{cm}^2/\text{s} \times 10^6$	°			
<b>G1:</b>	632	301	-1.2±0.08	7.3	13.5	0.9	9.3	15.8	50.0	18.3	341 <sup>1)</sup>			
			3.6±0.05	3.8	5.1		7.2	4.2	8.3	1.9				
	1035	412	-1.0±0.05	6.4	6.5	-1.6	63.9	11.9	26.2	6.7	333			
			2.1±0.05	2.3	6.3		2.7	19.1	41.3	10.3				
	1838	387	-0.7±0.06	6.1	8.7	-6.1	11.5	14.9	38.2	11.3	346			
			2.9±0.08	3.0	14.2		4.6	20.6	67.0	25.2				
<b>G2:</b>	370	168	-0.1±0.17	7.8	31.9	-3.9	23.2	17.0	83.0	46.9	355			
			1.4±0.12	1.4	14.6		1.0	7.7	25.5	9.7				
	977	240	0.8±0.08	5.5	9.1	-2.6	87.3	12.5	32.7	9.9	42			
			0.8±0.07	1.1	8.3		64.2	6.0	14.9	4.3				
<b>G3:</b>	367	18	8.0±0.10	11.0	6.7	0.9	14.0	1.1	2.4	0.6	128			
			-6.2±0.17	10.1	21.2		51.5	2.0	8.0	3.7				
	579	18	2.2±0.23	9.9	36.2		18.5	29.0	2.7	14.1	159			
			-5.9±0.18	6.3	21.8		19.6	1.8	7.1	3.3				
	881	18	2.0±0.12	6.7	9.9	2.7	7.6	2.8	7.6	2.4	145			
			-2.8±0.08	3.5	5.2		6.0	1.5	3.0	0.7				
	1383	18	1.3±0.07	4.5	3.3	-0.8	2.5	2.8	4.4	0.8	58			
			0.8±0.05	1.5	1.8		1.1	2.0	2.4	0.3				
	2085	18	0.1±0.08	4.2	2.4	0.8	3.0	2.6	3.4	0.5	3			
			1.4±0.14	1.4	3.6		0.9	1.6	2.7	0.5				

<sup>1)</sup> Rotor blocked from 15.4.94 - 5.6.94. Statistics calculated from 17.3.93 - 15.4.94.

Current statistics (low-passed data)

Id	sampl. depth	rec. length	$\bar{U}$ $V$	mag <b>v</b>	$\frac{\bar{U}'\bar{U}'}{V'V'}$		$\frac{\bar{U}'\bar{V}'}{V'V'}$		$k_E$ $K_M$	$T_x$ $T_y$	$L_x$ $L_y$	$K_x$ $K_y$	dir
					m	days	cm/s	cm/s	cm <sup>2</sup> /s <sup>2</sup>				
<b>G4:</b>	435	22	$1.9 \pm 0.09$	11.6	6.7	1.2	5.0	2.3	5.2	1.3	104		
			$-0.5 \pm 0.06$	2.0	3.2	2.0	2.0	2.1	3.1	0.5			
	587	22	$1.5 \pm 0.06$	8.1	3.5	-0.7	2.1	2.2	3.5	0.7	108		
			$-0.5 \pm 0.03$	1.6	0.7	1.3	1.3	1.7	1.2	0.1			
	739	22	$0.2 \pm 0.08$	7.1	5.5	-1.0	3.3	2.2	4.5	1.1	18		
			$0.6 \pm 0.04$	0.7	1.1	0.2	1.0	0.9	<0.1				
	941	22	$-1.8 \pm 0.16$	8.5	22.3	3.2	12.2	1.8	7.5	3.5	290		
			$0.7 \pm 0.05$	1.9	2.2	2.2	1.9	1.6	2.1	0.3			
	2485	22	$-3.8 \pm 0.10$	12.0	9.0	5.5	8.2	2.0	5.1	1.5	235		
			$-2.7 \pm 0.10$	4.7	7.5	10.9	3.8	9.1	2.5				
<b>G5:</b>	140	13	$-10.9 \pm 0.06$	12.4	3.2	-3.3	9.2	0.7	1.2	0.2	280		
			$1.9 \pm 0.12$	11.1	15.1		61.4	1.8	6.2	2.4			
	340	13	$-11.8 \pm 0.07$	14.0	5.0	-9.0	13.4	1.2	2.3	0.5	284		
			$2.9 \pm 0.15$	12.1	21.7		73.6	1.8	7.3	3.4			
	540	13	$-9.3 \pm 0.06$	13.1	3.1	-3.9	6.1	1.1	1.7	0.3	283		
			$2.1 \pm 0.10$	9.6	9.0		46.0	1.7	4.5	1.3			
	1140	13	$-1.9 \pm 0.03$	5.2	1.0	-0.5	1.1	0.8	0.7	<0.1	287		
			$0.6 \pm 0.03$	2.0	1.2		2.0	0.6	0.5	<0.1			
	2140	13	$-0.0 \pm 0.02$	3.2	0.5	-0.1	0.6	1.3	0.8	<0.1	349		
			$0.2 \pm 0.03$	0.2	0.7		<0.1	0.6	0.4	<0.1			






**A**a95047; sd= 632 m; 16. 6.1993 22:00 – 15. 4.1994 0:00 (UTC)

**B**a95048; sd= 1035 m; 16. 6.1993 22:00 – 5. 8.1994 14:00 (UTC)

**C**a95049; sd= 1838 m; 16. 6.1993 22:00 – 11. 7.1994 7:00 (UTC)


G1, record no. = 095047, sampling depth = 632 m





G1

record no: a95047, sampling interval = 60.0 minutes

u-component (zonal)




v-component (meridional)



16. 6.1993 22:00 – 15. 4.1994 0:00 (UTC)

Time marks (x) are given every 720 hours


G1, record no. = 095048, sampling depth = 1035 m





G1

record no: a95048, sampling interval = 60.0 minutes

u-component (zonal)




v-component (meridional)



16. 6.1993 22:00 – 5. 8.1994 14:00 (UTC)

Time marks (x) are given every 720 hours


G1, record no. = 095049, sampling depth = 1838 m





G1

record no: a95049, sampling interval = 60.0 minutes

u-component (zonal)




v-component (meridional)



16. 6.1993 22:00 – 11. 7.1994 7:00 (UTC)


Time marks (x) are given every 720 hours





**A**o95051; sd= 370 m; 17. 6.1993 13:00 – 4.12.1993 4:00 (UTC)  
**B**o95053; sd= 977 m; 17. 6.1993 13:00 – 15. 2.1994 2:00 (UTC)


G2, record no. = 095051, sampling depth = 370 m





G2

record no: a95051, sampling interval = 60.0 minutes

u-component (zonal)




v-component (meridional)



17. 6.1993 12:00 – 4.12.1993 4:00 (UTC)

Time marks (x) are given every 720 hours


G2, record no. = 095053, sampling depth = 977 m





G2

record no: a95053, sampling interval = 60.0 minutes

u-component (zonal)




v-component (meridional)



17. 6.1993 13:00 – 15. 2.1994 2:00 (UTC)

Time marks (x) are given every 720 hours






Time marks (x) are given every 48 hours

<b>A</b>	94037; sd= 367 m; 13. 6.1994	9:00 -	3. 7.1994	5:50 (UTC)
<b>B</b>	94038; sd= 579 m; 13. 6.1994	9:00 -	3. 7.1994	5:50 (UTC)
<b>C</b>	94039; sd= 881 m; 13. 6.1994	9:00 -	3. 7.1994	5:50 (UTC)
<b>D</b>	94040; sd= 1383 m; 13. 6.1994	9:00 -	3. 7.1994	5:50 (UTC)
<b>E</b>	94041; sd= 2085 m; 13. 6.1994	9:00 -	3. 7.1994	5:50 (UTC)


G3, record no. = 094037, sampling depth = 367 m





G3

record no: a94037, sampling interval = 10.0 minutes

u-component (zonal)




v-component (meridional)



13. 6.1994 9:00 – 3. 7.1994 5:50 (UTC)

Time marks (x) are given every 48 hours


G3, record no. = a94038, sampling depth = 579 m





G3

record no: a94038, sampling interval = 10.0 minutes

u-component (zonal)




v-component (meridional)



13. 6.1994 9:00 – 3. 7.1994 5:50 (UTC)

Time marks (x) are given every 48 hours


G3, record no. = 094039, sampling depth = 881 m





G3

record no: a94039, sampling interval = 10.0 minutes

u-component (zonal)




v-component (meridional)



13. 6.1994 9:00 – 3. 7.1994 5:50 (UTC)

Time marks (x) are given every 48 hours


G3, record no. = a94040, sampling depth = 1383 m





G3

record no: a94040, sampling interval = 10.0 minutes

u-component (zonal)




v-component (meridional)



13. 6.1994 9:00 – 3. 7.1994 5:50 (UTC)

Time marks (x) are given every 48 hours


G3, record no. = a94041, sampling depth = 2085 m





G3

record no: a94041, sampling interval = 10.0 minutes

u-component (zonal)




v-component (meridional)



13. 6.1994 9:00 – 3. 7.1994 5:50 (UTC)

Time marks (x) are given every 48 hours






Time marks (x) are given every 48 hours

- A** a94036; sd= 435 m; 14. 6.1994 12:20 – 8. 7.1994 7:50 (UTC)
- B** a94035; sd= 587 m; 14. 6.1994 12:20 – 8. 7.1994 7:50 (UTC)
- C** a94034; sd= 739 m; 14. 6.1994 12:20 – 8. 7.1994 7:50 (UTC)
- D** a94033; sd= 941 m; 14. 6.1994 12:20 – 8. 7.1994 7:50 (UTC)
- E** a94031; sd= 2485 m; 14. 6.1994 12:20 – 8. 7.1994 7:50 (UTC)


G4, record no. = a94036, sampling depth = 435 m





G4

record no: a94036, sampling interval = 10.0 minutes

u-component (zonal)




v-component (meridional)



14. 6.1994 12:20 – 8. 7.1994 7:50 (UTC)

Time marks (x) are given every 48 hours


G4, record no. = a94035, sampling depth = 587 m





G4

record no: a94035, sampling interval = 10.0 minutes

u-component (zonal)




v-component (meridional)



14. 6.1994 12:20 – 8. 7.1994 7:50 (UTC)



Time marks (x) are given every 48 hours

G4, record no. = a94034, sampling depth = 739 m



G4


record no: a94034, sampling interval = 10.0 minutes



14. 6.1994 12:20 – 8. 7.1994 7:50 (UTC)

Time marks (x) are given every 48 hours


G4, record no. = a94033, sampling depth = 941 m





G4

record no: a94033, sampling interval = 10.0 minutes

u-component (zonal)




v-component (meridional)



14. 6.1994 12:20 – 8. 7.1994 7:50 (UTC)

Time marks (x) are given every 48 hours


G4, record no. = a94031, sampling depth = 2485 m





G4

record no: a94031, sampling interval = 10.0 minutes

u-component (zonal)




v-component (meridional)



14. 6.1994 12:20 – 8. 7.1994 7:50 (UTC)

Time marks (x) are given every 48 hours






Time marks (x) are given every 48 hours

- A**a95036; sd= 140 m; 6. 8.1995 15:15 – 21. 8.1995 14:10 (UTC)
- B**a95037; sd= 340 m; 6. 8.1995 15:15 – 21. 8.1995 14:10 (UTC)
- C**a95038; sd= 540 m; 6. 8.1995 15:15 – 21. 8.1995 14:15 (UTC)
- D**a95039; sd= 1140 m; 6. 8.1995 15:15 – 21. 8.1995 14:10 (UTC)
- E**a95040; sd= 2140 m; 6. 8.1995 15:15 – 21. 8.1995 14:10 (UTC)


G5, record no. = 095036, sampling depth = 140 m





G5

record no: a95036, sampling interval = 5.0 minutes

u-component (zonal)




v-component (meridional)



6. 8.1995 15:15 – 21. 8.1995 14:10 (UTC)


Time marks (x) are given every 48 hours





G5

record no: a95037, sampling interval = 5.0 minutes

u-component (zonal)




v-component (meridional)



6. 8.1995 15:15 – 21. 8.1995 14:10 (UTC)

Time marks (x) are given every 48 hours


G5, record no. = 095038, sampling depth = 540 m





G5

record no: a95038, sampling interval = 5.0 minutes

u-component (zonal)




v-component (meridional)



6. 8.1995 15:15 – 21. 8.1995 14:15 (UTC)

Time marks (x) are given every 48 hours


G5, record no. = a95039, sampling depth = 1140 m





G5

record no: a95039, sampling interval = 5.0 minutes

u-component (zonal)




v-component (meridional)



6. 8.1995 15:15 – 21. 8.1995 14:10 (UTC)

Time marks (x) are given every 48 hours


G5, record no. = a95040, sampling depth = 2140 m





G5

record no: a95040, sampling interval = 5.0 minutes

u-component (zonal)



v-component (meridional)



6. 8.1995 15:15 – 21. 8.1995 14:10 (UTC)

Time marks (x) are given every 48 hours



L01




### Mooring Data L01

current meter	sampl. depth	mab	record no.	sampling time
-	m	m	-	UTC
<b>L01: 14.3.91 - 19.1.92, 42° 55.54'N, 9° 50.88'W, si = 60'</b>				
<u>RCM5 6966 435 1510 92008 14.3.91 14:00 - 10.1.92 08:00</u>				

mab = metres above bottom, si = sampling interval

Current statistics (low-passed data)


<i>Id</i>	samp1. depth	rec. length	$\bar{u}$	$\bar{v}$	mag $\bar{v}$	$\overline{u'u'}$ $\overline{v'v'}$	$\overline{u'v'}$	$k_E$ $K_N$	$T_x$ $T_y$	$L_x$ $L_y$	$K_x$ $K_y$	dir
m	days	cm/s	cm/s	cm/s	cm <sup>2</sup> /s <sup>2</sup>	cm <sup>2</sup> /s <sup>2</sup>	cm <sup>2</sup> /s <sup>2</sup>	days	km	cm <sup>2</sup> /s <sup>2</sup> × 10 <sup>6</sup>	°	
L01: 435	300	-0.9±0.08 1.9±0.13	9.7 2.0	13.7 30.4	4.4 2.1	22.1 2.1	7.1 5.5	22.7 26.4	8.4 14.6	335		



Time marks (x) are given every 720 hours

o92008; sd= 435 m; 14. 3.1991 14:00 – 10. 1.1992 8:00 (UTC)


L01, record no. = 0920008, sampling depth = 435 m




L01

record no: a92008, sampling interval = 60.0 minutes

u-component (zonal)



v-component (meridional)





AvH



### Mooring Data AvH


---

current meter	sampl. depth	mab	record no.	sampling time			
-	m	m	-	UTC			
<b>AvH: 1.10.89 - 9.8.90, 42° 32.80'N, 14° 28.60'W, si = 60'</b>							
RCM4	4248	410	4755	90143	2.10.89	04:00	-
RCM4	4364	1110	4055	90144	1.10.89	19:00	-
RCM5	6830	1810	3355	90145	1.10.89	19:00	-
RCM5	6863	3060	2105	90146	1.10.89	19:00	-
RCM5	6823	4640	524	90147	1.10.89	20:00	-
RCM5	6825	5045	120	90148	1.10.89	20:00	-
RCM5	6697	5144	21	90150	1.10.89	20:00	-
<u>RCM5</u>	<u>6839</u>	<u>5154</u>	<u>11</u>	<u>90151</u>	<u>1.10.89</u>	<u>22:00</u>	<u>-</u>
							<b>7.3.90 03:00</b>

mab = metres above bottom, si = sampling interval


Current statistics (low-passed data)

Id	sampler.	rec.	length	$\bar{u}$	$\bar{v}$	mag	$\overline{u'u'}$		$\overline{u'v'}$		$k_E$	$T_x$	$T_y$	$L_x$	$L_y$	$K_x$	$K_y$	dir
							$\overline{u'v'}$	$\overline{v'v'}$	$cm^2/s^2$	$cm^2/s^2$				days	km	$cm^2/s \times 10^6$	$^\circ$	
AVH:	410	308	-0.5±0.07	5.4	9.3	-0.8	9.1	13.0	34.5	10.5	210							
		-0.8±0.07	0.9	8.8	0.4		16.1	41.2	12.2									
	1110	306	-0.8±0.06	4.5	6.0	-0.2	5.4	17.6	37.3	9.2	207							
		-1.6±0.05	1.8	4.8		1.6	18.2	34.5	7.6									
	1810	309	-0.4±0.04	3.6	3.3	0.7	3.5	21.5	33.8	6.1	209							
		-0.7±0.04	0.8	3.6		0.3	16.0	26.4	5.0									
	3060	309	-0.2±0.04	3.2	3.1	0.7	3.6	24.4	37.3	6.6	225							
		-0.2±0.04	0.3	4.0		<0.1	15.0	25.9	5.2									
	4640	281	-0.3±0.06	4.3	6.3	0.5	7.0	22.3	48.5	12.2	230							
		-0.2±0.07	0.4	7.6		<0.1	15.0	35.8	9.9									
	5045	297	-0.2±0.05	3.4	4.5	1.0	4.6	20.5	37.7	8.0	225							
		-0.2±0.05	0.3	4.8		<0.1	16.0	30.2	6.6									
	5144	282	-0.2±0.06	4.3	6.0	0.6	5.9	20.7	43.8	10.8	207							
		-0.5±0.06	0.5	5.9		0.1	14.3	29.9	7.2									
	5154	154	-0.1±0.09	4.4	7.5	-0.4	8.4	22.5	53.3	14.6	258							
		-0.0±0.10	0.1	9.2		<0.1	15.5	40.6	12.3									



<b>A</b>	a90143; sd= 410 m;	2.10.1989	4:00 –	8. 8.1990	8:00 (UTC)
<b>B</b>	a90144; sd= 1110 m;	2.10.1989	4:00 –	5. 8.1990	16:00 (UTC)
<b>C</b>	a90145; sd= 1810 m;	2.10.1989	4:00 –	9. 8.1990	8:00 (UTC)
<b>D</b>	a90146; sd= 3060 m;	2.10.1989	4:00 –	9. 8.1990	8:00 (UTC)
<b>E</b>	a90147; sd= 4640 m;	2.10.1989	4:00 –	11. 7.1990	23:00 (UTC)
<b>F</b>	a90148; sd= 5045 m;	2.10.1989	4:00 –	27. 7.1990	13:00 (UTC)
<b>G</b>	a90150; sd= 5144 m;	2.10.1989	4:00 –	12. 7.1990	21:00 (UTC)
<b>H</b>	a90151; sd= 5154 m;	2.10.1989	4:00 –	7. 3.1990	3:00 (UTC)


AvH, record no. = 90143, sampling depth = 410 m





AvH

record no: a90143, sampling interval = 60.0 minutes

u-component (zonal)




v-component (meridional)



2.10.1989 4:00 – 8. 8.1990 8:00 (UTC)

Time marks (x) are given every 720 hours


AvH, record no. = a90144, sampling depth = 1110 m





AvH

record no: a90144, sampling interval = 60.0 minutes

u-component (zonal)




v-component (meridional)



1.10.1989 19:00 – 5. 8.1990 16:00 (UTC)

Time marks (x) are given every 720 hours


AvH, record no. = a90145, sampling depth = 1810 m





AvH

record no: a90145, sampling interval = 60.0 minutes

u-component (zonal)




v-component (meridional)



1.10.1989 19:00 – 9. 8.1990 8:00 (UTC)

Time marks (x) are given every 720 hours


AvH, record no. = a90146, sampling depth = 3060 m





AvH

record no: a90146, sampling interval = 60.0 minutes

u-component (zonal)




v-component (meridional)



1.10.1989 19:00 – 9. 8.1990 8:00 (UTC)

Time marks (x) are given every 720 hours


AvH, record no. = a90147, sampling depth = 4640 m





AvH

record no: a90147, sampling interval = 60.0 minutes

u-component (zonal)




v-component (meridional)



1.10.1989 20:00 – 11. 7.1990 23:00 (UTC)

Time marks (x) are given every 720 hours


AvH, record no. = 090148, sampling depth = 5045 m





AvH

record no: a90148, sampling interval = 60.0 minutes

u-component (zonal)




v-component (meridional)



1.10.1989 20:00 – 27. 7.1990 13:00 (UTC)

Time marks (x) are given every 720 hours


AvH, record no. = 990150, sampling depth = 5144 m





AvH

record no: a90150, sampling interval = 60.0 minutes

u-component (zonal)




v-component (meridional)



1.10.1989 20:00 – 12. 7.1990 21:00 (UTC)

Time marks (x) are given every 720 hours


AvH, record no. = 090151, sampling depth = 5154 m





AvH

record no: a90151, sampling interval = 60.0 minutes

u-component (zonal)



v-component (meridional)



1.10.1989 22:00 – 7. 3.1990 3:00 (UTC)

Time marks (x) are given every 720 hours



K1




### Mooring Data K1

current meter	sampl. depth	mab	record no.	sampling time
-	m	m	-	UTC
<b>K1: 19.3.91 - 13.4.91, 40° 47.4'N, 9° 06.2'W, si = 10'</b>				
RCM4	7242	40	65 91023	19.3.91 14:10 - 13.4.91 10:20
RCM4	8510	71	34 91024	19.3.91 14:10 - 13.4.91 08:50
RCM4	8509	98	7 91025	19.3.91 15:00 - 13.4.91 10:00

mab = metres above bottom, si = sampling interval

Current statistics (low-passed data)


Id	sampler. depth	rec. length	$\bar{U}$	$\bar{V}$	$mag$	$\frac{\bar{U}'\bar{U}'}{\bar{V}'\bar{V}'}$	$\bar{U}'\bar{V}'$	$k_E$	$T_x$	$L_x$	$K_x$	$dir$
			m	days	cm/s	cm/s	cm <sup>2</sup> /s <sup>2</sup>	cm <sup>2</sup> /s <sup>2</sup>	days	km	km	$L_y$
<b>K1:</b>	40	23	1.1±0.10	15.7	8.1	15.6	52.0	0.9	2.2	0.6	0.6	144
	71	23	-1.6±0.33	2.0	95.9		1.9	1.2	10.2	10.0		
		0.2±0.03	8.2	0.7	1.6	24.4	0.7	0.5	<0.1		173	
	98	23	-1.6±0.24	1.7	48.0		1.4	1.5	9.1	6.3		
		0.6±0.06	7.3	3.4	-9.8	23.9	1.1	1.8	0.3		148	
		-0.9±0.23	1.1	44.4		0.6	1.6	9.1	6.0			



Time marks (x) are given every 48 hours

**A**o91023; sd= 40 m; 19. 3.1991 15:00 – 13. 4.1991 10:20 (UTC)  
**B**o91024; sd= 71 m; 19. 3.1991 15:00 – 13. 4.1991 8:50 (UTC)  
**C**o91025; sd= 98 m; 19. 3.1991 15:00 – 13. 4.1991 10:00 (UTC)


K1, record no. = 091023, sampling depth = 40 m





K1

record no: a91023, sampling interval = 10.0 minutes

u-component (zonal)




v-component (meridional)



19. 3.1991 14:10 – 13. 4.1991 10:20 (UTC)

Time marks (x) are given every 48 hours


K1, record no. = 091024, sampling depth = 71 m





K1

record no: a91024, sampling interval = 10.0 minutes

u-component (zonal)




v-component (meridional)



19. 3.1991 14:10 – 13. 4.1991 8:50 (UTC)

Time marks (x) are given every 48 hours


K1, record no. = a91025, sampling depth = 98 m





K1

record no: a91025, sampling interval = 10.0 minutes

u-component (zonal)



v-component (meridional)



19. 3.1991 15:00 – 13. 4.1991 10:00 (UTC)

Time marks (x) are given every 48 hours



WH



### Mooring Data WH


current meter	sampl. depth	mab	record no.	sampling time
-	m	m	-	UTC
<b>WH: 15.8.90 - 27.8.91, 40° 13.3'N, 10° 15.7'W, si = 60'</b>				
RCM5	6787	1164	2736 91088	15.8.90 12:00 - 26.8.91 05:00 <sup>1)</sup>
RCM5	6785	1564	2336 91089	15.8.90 12:00 - 27.8.91 07:00

mab = metres above bottom, si = sampling interval

<sup>1)</sup> speed only, no direction

Current statistics (low-passed data)


$\overline{u}$	$\overline{v}$	$\overline{u'v'}$	$\overline{u'u'}$	$\overline{v'v'}$	$k_E$	$T_x$	$L_x$	$K_x$	$dir$	
$\overline{u}$	$\overline{v}$	$\frac{\overline{u'u'}}{\nabla' \nabla'}$	$\frac{\overline{u'u'}}{\nabla' \nabla'}$	$\frac{\overline{v'v'}}{\nabla' \nabla'}$	$k_M$	$T_y$	$L_y$	$K_y$		
m	days	cm/s	cm/s	cm <sup>2</sup> /s <sup>2</sup>	cm <sup>2</sup> /s <sup>2</sup>	cm <sup>2</sup> /s <sup>2</sup>	days	km	cm <sup>2</sup> /s <sup>2</sup> × 10 <sup>6</sup>	
WH: 1564	377	-0.8 ± 0.04 0.3 ± 0.04	3.0 0.8	3.0 3.4	-0.1 0.3	3.2 0.3	8.0 20.1	12.0 31.9	2.1 5.9	289



Time marks (x) are given every 720 hours

a91089; sd= 1564 m; 15. 8.1990 12:00 – 27. 8.1991 7:00 (UTC)


WH, record no. = a91089, sampling depth = 1564 m




WH

record no: a91089, sampling interval = 60.0 minutes

u-component (zonal)



v-component (meridional)





P3



### Mooring Data P3


---

current meter	sampl. depth	mab	record no.	sampling time
-	m	m	-	UTC
<b>P3: 30.7.83 - 14.1.84, 40° 1.8'N, 9° 45.2'W, si = 30'</b>				
RCM4	6559	260	650 83081	30.7.83 11:40 - 14.1.84 07:10
RCM4	5933	410	500 83080	30.7.83 11:40 - 14.1.84 07:40
RCM4	6143	650	260 83079	30.7.83 11:40 - 14.1.84 07:10
RCM4	4889	810	100 83078	30.7.83 11:40 - 14.1.84 07:10

mab = metres above bottom, si = sampling interval

Current statistics (low-passed data)


$\overline{u}$	$\overline{v}$	$mag$	$\overline{u'u'}$	$\overline{u'v'}$	$k_x$	$T_x$	$L_x$	$K_x$	$dir$	
m	days	cm/s	cm/s	cm <sup>2</sup> /s <sup>2</sup>	cm <sup>2</sup> /s <sup>2</sup>	days	km	cm <sup>2</sup> /s <sup>2</sup> x 10 <sup>6</sup>	°	
<b>P3:</b>	260	166	0.4±0.06	9.2	8.1	-2.1	23.9	5.2	12.9	3.7
			3.8±0.14	3.8	39.7		7.2	10.3	55.9	35.2
	410	166	0.0±0.05	7.9	4.7	0.8	18.9	2.8	5.2	1.1
			2.8±0.13	2.8	33.1		3.8	9.0	44.7	25.8
	650	166	0.5±0.03	6.9	1.4	1.3	10.8	1.6	1.7	0.2
			3.3±0.10	3.4	20.3		5.7	4.7	18.3	8.3
	810	166	0.7±0.02	7.6	1.0	-0.9	10.1	1.4	1.2	0.1
			3.6±0.10	3.6	19.2		6.7	3.3	12.6	5.5



Time marks (x) are given every 720 hours

- A**o83081; sd= 260 m; 30. 7.1983 11:40 – 14. 1.1984 7:10 (UTC)
- B**o83080; sd= 410 m; 30. 7.1983 11:40 – 14. 1.1984 7:40 (UTC)
- C**o83079; sd= 650 m; 30. 7.1983 11:40 – 14. 1.1984 7:10 (UTC)
- D**o83078; sd= 810 m; 30. 7.1983 11:40 – 14. 1.1984 7:10 (UTC)


P3, record no. = a83081, sampling depth = 260 m





P3

record no: a83081, sampling interval = 30.0 minutes

u-component (zonal)




v-component (meridional)



30. 7.1983 11:40 – 14. 1.1984 7:10 (UTC)



Time marks (x) are given every 720 hours

P3, record no. = a83080, sampling depth = 410 m



P3


record no: a83080, sampling interval = 30.0 minutes



30. 7.1983 11:40 – 14. 1.1984 7:40 (UTC)

Time marks (x) are given every 720 hours


P3, record no. = 083079, sampling depth = 650 m





P3

record no: a83079, sampling interval = 30.0 minutes

u-component (zonal)




v-component (meridional)



30. 7.1983 11:40 – 14. 1.1984 7:10 (UTC)

Time marks (x) are given every 720 hours


P3, record no. = 083078, sampling depth = 810 m





P3

record no: a83078, sampling interval = 30.0 minutes

u-component (zonal)



v-component (meridional)



30. 7.1983 11:40 – 14. 1.1984 7:10 (UTC)

Time marks (x) are given every 720 hours



CS



### Mooring Data CS


---

current meter	sampl. depth	mab	record no.	sampling time		
-	m	m	-	UTC		
<b>CS: 28.3.92 - 19.5.92, 36° 37.1'N, 9° 22.0'W, si = 60'</b>						
RCM8	10586	235	1725	92047	28.3.92	19:00 - 19.5.92 10:00
RCM5	6833	337	1623	92048	28.3.92	19:00 - 19.5.92 10:00
RCM8	10590	538	1422	92049	28.3.92	19:00 - 19.5.92 10:00
RCM8	10588	753	1207	92050	28.3.92	19:00 - 19.5.92 08:00
RCM5	6966	1254	706	92051	28.3.92	19:00 - 19.5.92 10:00
RCM8	10081	1855	105	92052	28.3.92	19:00 - 19.5.92 10:00

mab = metres above bottom, si = sampling interval

Current statistics (low-passed data)


<i>ID</i>	sampl.	rec.	$\bar{U}$	$\bar{V}$	mag	$\frac{\overline{U'U'}}{\overline{V'V'}}$	$\overline{U'V'}$	$k_E$	$T_x$	$L_x$	$K_x$	dir
	depth	length	cm/s	cm/s	cm/s	cm <sup>2</sup> /s <sup>2</sup>	cm <sup>2</sup> /s <sup>2</sup>	km	$T_y$	$L_y$	$K_y$	
<b>CS:</b>	235	50	-5.3±0.37	12.1	44.2	-12.4	47.6	3.0	17.5	11.6	241	
			-2.9±0.41	6.1	51.0		18.5	3.7	23.1	16.5		
	337	50	-5.2±0.23	10.2	17.1	-2.4	23.8	2.7	9.8	4.0	248	
			-2.1±0.31	5.6	30.4		15.6	3.4	16.0	8.8		
	538	50	-7.5±0.37	12.1	42.7	2.7	33.6	1.5	8.7	5.7	278	
			1.1±0.28	7.8	24.5		30.6	3.2	13.8	6.8		
	753	50	-10.2±0.37	15.1	41.6	9.5	55.2	1.8	9.9	6.4	296	
			4.9±0.47	11.3	68.9		63.6	6.1	44.0	36.5		
	1254	50	-0.1±0.41	10.8	51.9	-29.4	60.3	1.6	10.1	7.2	346	
			0.6±0.47	0.6	68.8		0.2	3.5	24.8	20.6		
	1855	50	-1.1±0.05	4.6	8.9	-1.6	4.9	0.8	2.1	0.6	289	
			0.4±0.05	1.2	0.8		0.7	0.8	0.6	<0.1		



Time marks (x) are given every 720 hours

- A**a92047; sd= 235 m; 28. 3.1992 19:00 – 19. 5.1992 10:00 (UTC)
- B**a92048; sd= 337 m; 28. 3.1992 19:00 – 19. 5.1992 10:00 (UTC)
- C**a92049; sd= 538 m; 28. 3.1992 19:00 – 19. 5.1992 10:00 (UTC)
- D**a92050; sd= 753 m; 28. 3.1992 19:00 – 19. 5.1992 8:00 (UTC)
- E**a92051; sd= 1254 m; 28. 3.1992 19:00 – 19. 5.1992 10:00 (UTC)
- F**a92052; sd= 1855 m; 28. 3.1992 19:00 – 19. 5.1992 10:00 (UTC)


CS, record no. = a92047, sampling depth = 235 m





CS

record no: a92047, sampling interval = 60.0 minutes

u-component (zonal)




v-component (meridional)



28. 3.1992 19:00 – 19. 5.1992 10:00 (UTC)

Time marks (x) are given every 240 hours


CS, record no. = a92048, sampling depth = 337 m





CS

record no: a92048, sampling interval = 60.0 minutes

u-component (zonal)




v-component (meridional)



28. 3.1992 19:00 – 19. 5.1992 10:00 (UTC)



Time marks (x) are given every 240 hours

CS, record no. = a92049, sampling depth = 538 m



CS


record no: a92049, sampling interval = 60.0 minutes



28. 3.1992 19:00 – 19. 5.1992 10:00 (UTC)



Time marks (x) are given every 240 hours

CS, record no. = a92050, sampling depth = 753 m



CS


record no: a92050, sampling interval = 60.0 minutes



28. 3.1992 19:00 – 19. 5.1992 8:00 (UTC)

Time marks (x) are given every 240 hours


CS, record no. = a92051, sampling depth = 1254 m





CS

record no: a92051, sampling interval = 60.0 minutes

u-component (zonal)




v-component (meridional)



28. 3.1992 19:00 – 19. 5.1992 10:00 (UTC)



Time marks (x) are given every 240 hours

CS, record no. = a92052, sampling depth = 1855 m



CS

record no: a92052, sampling interval = 60.0 minutes



28. 3.1992 19:00 – 19. 5.1992 10:00 (UTC)

Time marks (x) are given every 240 hours



# AL1, AL2



### Mooring Data AL1 and AL2


---

current meter	sampl. depth	mab	record no.	sampling time			
-	m	m	-	UTC			
<b>AL1: 28.10.91 - 14.1.92, 36° 14.16'N, 8° 36.12'W, si = 30'</b>							
RCM4	8509	358	1678	92001	20.10.91	12:30	- 14.1.92 09:00
RCM4	7242	510	1526	92002	20.10.91	12:30	- 14.1.92 09:00
RCM7	9750	811	1225	92003	20.10.91	12:30	- 14.1.92 09:00
RCM8	9386	1214	822	92004	20.10.91	12:30	- 14.1.92 09:00
RCM8	10081	1715	321	92005	20.10.91	12:30	- 14.1.92 09:00
RCM8	10082	2016	20	92006	20.10.91	12:30	- 14.1.92 09:00
<b>AL2: 14.1.92 - 19.5.92, 36° 14.20'N, 8° 35.10'W, si = 30'</b>							
RCM5	6786	309	1689	92041	14.1.92	15:00	- 19.5.92 04:30
RCM5	6703	471	1527	92042	14.1.92	15:00	- 19.5.92 05:00
RCM5	6701	773	1255	92043	14.1.92	15:00	- 19.5.92 05:00
RCM5	6700	1176	822	92044	14.1.92	15:00	- 19.5.92 05:30
RCM4	6129	1677	321	92045	14.1.92	15:00	- 19.5.92 05:30
RCM5	6119	1978	20	92046	14.1.92	15:00	- 27.3.92 16:00

mab = metres above bottom, si = sampling interval

Current statistics (low-passed data)


Id	sampl. depth	rec. length	$\bar{u}$ $\bar{v}$	mag $\mathbf{v}$	$\overline{u'u'}$		$\overline{u'v'}$		$k_E$ $k_W$	$T_x$ $T_y$	$L_x$ $L_y$	$K_x$ $K_y$	dir °
					m	days	cm/s	cm/s					
<b>All:</b>	358	84	4.5±0.13	9.3	17.3	-6.5	13.4	9.7	34.9	14.5	43		
			4.8±0.09	6.5	9.5	21.5	3.9	10.3	3.2				
	510	84	3.1±0.10	6.6	11.6	-2.6	8.3	9.9	29.1	9.9	53		
			2.4±0.07	3.9	4.9	7.6	3.5	6.6	1.5				
	811	84	4.5±0.18	9.6	33.3	-3.5	32.4	7.7	38.6	22.3	105		
			-1.2±0.17	4.6	31.6	10.7	3.8	18.3	10.3				
	1214	84	3.5±0.26	13.9	70.1	2.4	98.6	5.7	41.2	34.5	136		
			-3.7±0.35	5.1	127.1	2.4	13.3	4.4	43.3	48.9			
	1750	84	0.7±0.05	7.4	3.0	2.3	5.7	2.9	4.3	7.5	144		
			-0.9±0.09	1.1	8.3	0.1	0.6	4.7	11.8	3.4			
	2016	84	1.1±0.04	4.0	1.4	-0.1	1.7	1.7	1.7	0.2	72		
			0.4±0.04	1.2	2.1	0.1	0.7	2.7	3.4	0.5			
<b>All2:</b>	309	124	2.3±0.14	8.6	29.7	-1.7	23.2	13.1	62.0	33.8	28		
			4.3±0.10	4.8	16.7	11.7	8.8	31.0	12.6				
	471	124	1.4±0.12	6.1	23.1	2.4	17.1	10.0	41.6	20.0	31		
			2.3±0.08	2.7	11.2	3.6	8.5	24.5	8.2				
	773	124	1.7±0.12	8.5	23.0	-1.2	29.2	8.1	33.7	16.2	25		
			3.6±0.15	3.9	35.5	7.7	5.8	29.7	17.7				
	1176	124	-0.1±0.03	8.4	1.1	-0.3	1.0	0.7	0.6	<0.1	335		
			0.2±0.02	0.2	1.0	<0.1	0.7	0.6	<0.1				
	1677	124	-0.2±0.04	4.4	2.5	-1.3	4.1	2.4	3.2	0.5	346		
			1.0±0.06	1.0	5.7	<0.1	4.5	9.3	2.2				
	1978	71	0.7±0.03	3.9	0.6	0.6	2.9	1.1	0.7	<0.1	22		
			1.8±0.08	1.9	5.2	1.9	2.0	4.0	0.9				



Time marks (x) are given every 720 hours



- A**92001; sd= 358 m; 20.10.1991 12:30 – 14. 1.1992 9:00 (UTC)
- B**92002; sd= 510 m; 20.10.1991 12:30 – 14. 1.1992 9:00 (UTC)
- C**92003; sd= 811 m; 20.10.1991 12:30 – 14. 1.1992 9:00 (UTC)
- D**92004; sd= 1214 m; 20.10.1991 12:30 – 14. 1.1992 9:00 (UTC)
- E**92005; sd= 1750 m; 20.10.1991 12:30 – 14. 1.1992 9:00 (UTC)
- F**92006; sd= 2016 m; 20.10.1991 12:30 – 14. 1.1992 9:00 (UTC)

AL1, record no. = 092001, sampling depth = 358 m



AL1


record no: a92001, sampling interval = 30.0 minutes



20.10.1991 12:30 – 14. 1.1992 9:00 (UTC)

Time marks (x) are given every 720 hours


AL1, record no. = a920002, sampling depth = 510 m





AL1

record no: a92002, sampling interval = 30.0 minutes

u-component (zonal)




v-component (meridional)



20.10.1991 12:30 – 14. 1.1992 9:00 (UTC)

Time marks (x) are given every 720 hours

AL1, record no. = a92003, sampling depth = 811 m





AL1

record no: a92003, sampling interval = 30.0 minutes

u-component (zonal)


v-component (meridional)



20.10.1991 12:30 – 14. 1.1992 9:00 (UTC)

Time marks (x) are given every 720 hours


AL1, record no. = a92004, sampling depth = 1214 m





AL1

record no: a92004, sampling interval = 30.0 minutes

u-component (zonal)




v-component (meridional)



20.10.1991 12:30 – 14. 1.1992 9:00 (UTC)



Time marks (x) are given every 720 hours

AL1, record no. = 092005, sampling depth = 1750 m



AL1


record no: a92005, sampling interval = 30.0 minutes



20.10.1991 12:30 – 14. 1.1992 9:00 (UTC)

Time marks (x) are given every 720 hours


AL1, record no. = 092006, sampling depth = 2016 m





AL1

record no: a92006, sampling interval = 30.0 minutes

u-component (zonal)




v-component (meridional)



20.10.1991 12:30 – 14. 1.1992 9:00 (UTC)

Time marks (x) are given every 720 hours






Time marks (x) are given every 720 hours

Ao92041; sd=	309 m; 14. 1.1992	15:00 – 19. 5.1992	4:00 (UTC)
Ba92042; sd=	471 m; 14. 1.1992	15:00 – 19. 5.1992	5:00 (UTC)
Co92043; sd=	773 m; 14. 1.1992	15:00 – 19. 5.1992	5:00 (UTC)
Do92044; sd=	1176 m; 14. 1.1992	15:00 – 19. 5.1992	5:30 (UTC)
Eo92045; sd=	1677 m; 14. 1.1992	15:00 – 19. 5.1992	5:30 (UTC)
Fo92046; sd=	1978 m; 14. 1.1992	15:00 – 27. 3.1992	16:00 (UTC)


AL2, record no. = a92041, sampling depth = 309 m





AL2

record no: a92041, sampling interval = 30.0 minutes

u-component (zonal)




v-component (meridional)



14. 1.1992 15:00 – 19. 5.1992 4:00 (UTC)

Time marks (x) are given every 720 hours


AL2, record no. = a92042, sampling depth = 471 m





AL2

record no: a92042, sampling interval = 30.0 minutes

u-component (zonal)




v-component (meridional)



14. 1.1992 15:00 – 19. 5.1992 5:00 (UTC)



Time marks (x) are given every 720 hours

AL2, record no. = 092043, sampling depth = 773 m



AL2


record no: a92043, sampling interval = 30.0 minutes



14. 1.1992 15:00 – 19. 5.1992 5:00 (UTC)

Time marks (x) are given every 720 hours


AL2, record no. = 092044, sampling depth = 1176 m





AL2

record no: a92044, sampling interval = 30.0 minutes

u-component (zonal)




v-component (meridional)



14. 1.1992 15:00 – 19. 5.1992 5:30 (UTC)

Time marks (x) are given every 720 hours


AL2, record no. = 092045, sampling depth = 1677 m





AL2

record no: a92045, sampling interval = 30.0 minutes

u-component (zonal)




v-component (meridional)



14. 1.1992 15:00 – 19. 5.1992 5:30 (UTC)



Time marks (x) are given every 720 hours

AL2, record no. = a92046, sampling depth = 1978 m



AL2

record no: a92046, sampling interval = 30.0 minutes



14. 1.1992 15:00 – 27. 3.1992 16:00 (UTC)

Time marks (x) are given every 720 hours



R



### Mooring Data R

---


current meter	sampl. depth	mab	record no.	sampling time
-	m	m	-	UTC
<b>R: 20.10.91 - 15.4.92, 34° 29.1'N, 7° 50.8'W, si = 30'</b>				
RCM7	9084	538	1457	92036 19.10.91 12:00 - 16.4.92 16:30
RCM7	9053	791	1204	92037 19.10.91 12:00 - 18.5.92 18:30
RCM8	9754	1192	803	92038 19.10.91 12:00 - 7.3.92 19:00
RCM8	10079	1693	302	92039 19.10.91 12:00 - 1.3.92 07:00
RCM8	10080	1975	20	92040 19.10.91 12:00 - 22.3.92 18:00

mab = metres above bottom, si = sampling interval

Current statistics (low-passed data)

	Id	samp1.	rec.	$\bar{u}$	$\bar{v}$	mag $\bar{v}$	$\frac{\bar{u}'\bar{u}'}{\bar{v}'\bar{v}'}$	$\frac{\bar{u}'\bar{v}'}{\bar{v}'\bar{v}'}$	$K_E$ $K_M$	$T_x$ $T_y$	$L_x$ $L_y$	$K_x$ $K_y$	dir
	m	days	cm/s	cm/s	cm/s	cm <sup>2</sup> /s <sup>2</sup>	cm <sup>2</sup> /s <sup>2</sup>	cm <sup>2</sup> /s <sup>2</sup>	days	km	cm <sup>2</sup> /s <sup>2</sup> × 10 <sup>6</sup>	°	
R:	538	178	3.1±0.06	6.2	8.7	0.7	8.0	19.9	50.8	15.0	59		
	791	210	1.9±0.06	3.6	7.3	6.6	7.3	16.9	4.6				
	1192	138	2.1±0.04	5.9	5.4	2.0	6.7	13.7	27.4	6.4	45		
	1693	132	2.1±0.05	3.0	8.0	4.5	19.6	48.0	13.6				
	1975	91	-0.1±0.03	3.6	1.5	0.5	1.8	4.0	4.3	0.5	191		
			-0.3±0.03	0.3	2.2	<0.1	9.4	11.9	1.7				
			0.1±0.05	4.4	3.8	2.4	4.1	8.8	14.8	2.9	123		
			-0.0±0.05	0.1	4.4	<0.1	11.5	20.7	4.3				
			2.2±0.06	6.7	4.7	2.4	3.6	2.9	5.4	1.2	42 <sup>1)</sup>		
			2.5±0.05	3.3	2.6	5.4	3.1	4.3	0.7				


<sup>1)</sup> Rotor blocked from 19.1.92 - 5.3.92. Statistics calculated from 20.10.91 - 19.1.92.



Time marks (x) are given every 720 hours

- Aa92036;** sd= 538 m; 19.10.1991 12:00 – 16. 4.1992 16:30 (UTC)
- Ba92037;** sd= 791 m; 19.10.1991 12:00 – 18. 5.1992 18:30 (UTC)
- Ca92038;** sd= 1192 m; 19.10.1991 12:00 – 7. 3.1992 19:00 (UTC)
- Da92039;** sd= 1693 m; 19.10.1991 12:00 – 1. 2.1992 7:00 (UTC)
- Ea92040;** sd= 1975 m; 19.10.1991 12:00 – 19. 1.1992 10:00 (UTC)


R, record no. = a92036, sampling depth = 538 m





R

record no: a92036, sampling interval = 30.0 minutes

u-component (zonal)




v-component (meridional)



19.10.1991 12:00 – 16. 4.1992 16:30 (UTC)

Time marks (x) are given every 720 hours


R, record no. = a92037, sampling depth = 791 m





R

record no: a92037, sampling interval = 30.0 minutes

u-component (zonal)




v-component (meridional)



19.10.1991 12:00 – 18. 5.1992 18:30 (UTC)

Time marks (x) are given every 720 hours


R, record no. = a92038, sampling depth = 1192 m





R

record no: a92038, sampling interval = 30.0 minutes

u-component (zonal)




v-component (meridional)



19.10.1991 12:00 – 7. 3.1992 19:00 (UTC)

Time marks (x) are given every 720 hours


R, record no. = 092039, sampling depth = 1693 m





R

record no: a92039, sampling interval = 30.0 minutes

u-component (zonal)




v-component (meridional)



19.10.1991 12:00 – 1. 2.1992 7:00 (UTC)

Time marks (x) are given every 720 hours


R, record no. = a92040, sampling depth = 1975 m





R

record no: a92040, sampling interval = 30.0 minutes

u-component (zonal)



v-component (meridional)



19.10.1991 12:00 – 19. 1.1992 10:00 (UTC)

Time marks (x) are given every 720 hours



# M3




### Mooring Data M3

current meter	sampl. depth	mab	record no.	sampling time
-	m	m	-	UTC
<b>M3: 18.1.92 - 25.5.92, 32° 35.72'N, 10° 49.18'W, si = 30'</b>				
RCM8	10086	169	3810	92057 18.1.92 10:30 - 25.5.92 13:00

mab = metres above bottom, si = sampling interval


Current statistics (low-passed data)

$\overline{u}$	$\overline{v}$	$mag$	$\overline{u'u'}$	$\overline{u'v'}$	$k_E$	$T_x$	$L_x$	$K_x$	$dir$	
m	days	cm/s	cm/s	cm <sup>2</sup> /s <sup>2</sup>	cm <sup>2</sup> /s <sup>2</sup>	days	km	cm <sup>2</sup> /s <sup>2</sup> × 10 <sup>6</sup>	°	
M3:	169	124	-0.9 ± 0.14 -6.4 ± 0.21	11.7 6.5	29.5 66.5	16.7 21.0	48.0 17.4	6.9 122.3	32.3 99.8	17.6 188



a92057; sd= 169 m; 18. 1.1992 10:30 – 23. 5.1992 13:00 (UTC)


M3, record no. = a92057, sampling depth = 169 m




M3

record no: a92057, sampling interval = 30.0 minutes

u-component (zonal)



v-component (meridional)





## Berichte des Bundesamtes für Seeschiffahrt und Hydrographie

### Verzeichnis der veröffentlichten Arbeiten

- 
- 1 (1994) Sy, A., Ulrich, J. North Atlantic Ship-of-Opportunity XBT Programme 1990 - Data Report, 134 pp.
- 2 (1994) Hagen, E. Mittelstaedt, E. Feistel, R., Klein, H. Hydrographische Untersuchungen im Ostrandstromsystem vor Portugal und Marokko 1991 - 1992, 49 pp.
- 3 (1994) Oliczewski, J. Schmidt, D. Entwicklung einer Bestrahlungsapparatur zum photochemischen Aufschluß von Meerwasserproben zur Bestimmung von Schwermetallen, 70 pp.
- 4 (1994) BSH [Hrsg.] Das UN-Seerechtsübereinkommen tritt in Kraft: Inhalte und Konsequenzen für die Bundesrepublik Deutschland, 71 pp.
- 5 (1995) BSH [Hrsg.] Nationale Folgerungen aus dem Inkrafttreten des UN-Seerechtsübereinkommens, 103 pp.
- 6 (1995) Haffer, E. Schmidt, D. Entwicklung eines Probenvorbereitungsverfahrens zur Bestimmung von Arsen im Meerwasser mit der Totalreflexions-Röntgenfluoreszenzanalyse, 109 pp.
- 7 (1995) BSH [Hrsg.] Global Ocean Observing System - Statusbericht, 100 pp.
- 8 (1996) Mittelstaedt, E. Meincke, J., Klein, H. WOCE-Current measurements: The ACM8 array – Data Report, 150 pp.
- 9 (1996) BSH [Hrsg] GOOS Workshop - Anforderungen an ein wissenschaftliches Konzept für den deutschen Beitrag, 60 pp.
- 10 (1997) Sterzenbach, D. Entwicklung eines Analyseverfahrens zur Bestimmung von chlorierten Kohlenwasserstoffen in marinen Sedimenten und Schwebstoffen unter besonderer Berücksichtigung der überkritischen Fluidextraktion, 233 pp.
- 11 (1997) Jonas, M. Richter, R. Stand und Entwicklungstendenzen nautischer Systeme, Anlagen und Geräte an Bord von Seeschiffen, 37 pp.
- 12 (1997) Wedekind, C., Gabriel, H., Goroncy, I., Främke, G., Kautsky, H. "Meteor"-Reise Nr. 71/1985, Norwegen-Grönlandsee – Datenbericht. 44 pp.
- 13 (1998) BSH [Hrsg] HELCOM Scientific Workshop - The Effects of the 1997 Flood of the Odra and Vistula Rivers. 46 pp.
- 14 (1998) Berger, R., Klein, H., Mittelstaedt, E., Ricklefs, K., Ross, J. Der Wasseraustausch im Tidebecken Hörnum-Tief – Datenreport. 260 pp.
- 15 (1998) Röske, F. Wasserstandsvorhersage mittels neuronaler Netze. 212 pp.
- 16 (1998) Ross, J., Mittelstaedt, E., Klein, H., Berger, R., Ricklefs, K. Der Wasseraustausch im Tidebecken Hörnum-Tief – Abschlußbericht. 98 pp.

- 17 (1998) Klein, H. OPUS-Current Measurements: Mecklenburg Bight and Fehmarnbelt - Data Report, 150 pp
- 18 (1999) BSH [Hrsg] Deutscher Programmbeitrag zum Globalen Ozeanbeobachtungssystem (GOOS), 67 pp.
- 19 (1999) BSH [Hrsg] German Programme Contribution to the Global Ocean Observing System (GOOS), 71 pp.
- 20 (1999) Sztobryn, M. Ice Conditions in the Szczecin and Pomeranian Bay During the Normal Period from 1961-1990, 36 pp.  
Stanislawczyk, I  
Schmelzer, N.
- 21 (1999) Nies, H., M. Karcher Transportmechanismen radioaktiver Substanzen im Arktischen Ozean Bahe, C. Backhaus, J. - Numerische und experimentelle Studien am Beispiel der Barents- und Harms, I. 134 pp.
- 22 (2000) Lorbacher, K. Niederfrequente Variabilität meridionaler Transporte in der Divergenzzone des nordatlantischen Subtropen- und Subpolarwirbels – Der WOCE-Schnitt A2, 156 pp.
- 23 (2000) Klein, H. The Subsurface Eastern Boundary Current of the North Atlantic between 32°N and 53°N – Data Report, 240 pp.
- 24 (2000) Klein, H. Strömungen und Seegangsverhältnisse westlich der Insel Hiddensee – Datenreport, 59 pp.